arXiv:0804.0639v1 [math.RA] 4 Apr 2008

Gröbner-Shirshov bases for extensions of algebras*

Yuqun Chen

School of Mathematical Sciences
South China Normal University
Guangzhou 510631
P. R. China
yqchen@scnu.edu.cn

Abstract. An algebra $\mathcal{R}$ is called an extension of the algebra M by B if $M^2 = 0$, M is an ideal of $\mathcal{R}$ and $\mathcal{R}/M \cong B$ as algebras. In this paper, by using the Gröbner-Shirshov bases, we characterize completely the extensions of M by B. An algorithm to find the conditions of an algebra A to be an extension of M by B is obtained.

Keywords: algebra, module, Gröbner-Shirshov bases, extension
AMS 2000 Subject Classification: 16S15, 13P10

1 Preliminaries

Let k be a field, X a set, X^* the monoid of all words on X and X^+ the free semigroup of nonempty words on X. We denote $k\langle X_+\rangle$ the k-span of all nonempty words in X. As we know, $k\langle X_+\rangle$ is a free associative algebra without identity and $k\langle X\rangle$ a free associative algebra with identity. It is clear that, for every algebra $\mathscr{A}$ (not necessarily with 1), we have $\mathscr{A} \cong k\langle X_+\rangle/I$ for some X and ideal I of $k\langle X_+\rangle$. For a word $w \in X^*$, we denote the length of w by $deg(w)$. Let X^* be a well ordered set. Let $f \in k\langle X\rangle$ with the leading word $\bar{f}$. We say that f is monic if $\bar{f}$ has coefficient 1.

Definition 1.1 *([11], see also [2], [3]) Let f and g be two monic polynomials in $k\langle X\rangle$ and $<$ a well order on X^*. Then, there are two kinds of compositions:*

(1) *If w is a word such that $w = \bar{f}b = a\bar{g}$ for some $a, b \in X^*$ with $deg(\bar{f}) + deg(\bar{g}) > deg(w)$, then the polynomial $(f, g)_w = fb - ag$ is called the intersection composition of f and g with respect to w.*

(2) *If $w = \bar{f} = a\bar{g}b$ for some $a, b \in X^*$, then the polynomial $(f, g)_w = f - agb$ is called the inclusion composition of f and g with respect to w.*

Definition 1.2 *([2], [3], cf. [11]) Let $S \subseteq k\langle X\rangle$ with each $s \in S$ monic. Then the composition $(f, g)_w$ is called trivial modulo S if $(f, g)_w = \sum \alpha_i a_i s_i b_i$, where each $\alpha_i \in k$,*

---

*The research is supported by the National Natural Science Foundation of China (Grant No.10771077) and the Natural Science Foundation of Guangdong Province (Grant No.06025062).

$a_i, b_i \in X^*$ and $\overline{a_i s_i b_i} < w$. If this is the case, then we write

$$(f, g)_w \equiv 0 \quad mod(S, w)$$

In general, for $p, q \in k\langle X\rangle$, we write

$$p \equiv q \quad mod(S, w)$$

which means that $p - q = \sum \alpha_i a_i s_i b_i$, where each $\alpha_i \in k, a_i, b_i \in X^*$ and $\overline{a_i s_i b_i} < w$.

Definition 1.3 *([2], [3], cf. [11]) We call the set S with respect to the well order a Gröbner-Shirshov set (basis) in $k\langle X\rangle$ if any composition of polynomials in S is trivial relative to S.*

A well order $>$ on X^* is monomial if it is compatible with the multiplication of words, that is, for $u, v \in X^*$, we have

$$u > v \Rightarrow w_1 u w_2 > w_1 v w_2, \ for\ all\ w_1,\ w_2 \in X^*.$$

A standard example of monomial order on X^* is the deg-lex order to compare two words first by degree and then lexicographically, where X is a linearly ordered set.

The following lemma was proved by Shirshov [11] for the free Lie algebras (with deg-lex ordering) in 1962 (see also Bokut [2]). In 1976, Bokut [3] specialized the approach of Shirshov to associative algebras (see also Bergman [1]). For commutative polynomials, this lemma is known as the Buchberger's Theorem in [5] and [6].

Lemma 1.4 *(Composition-Diamond Lemma) Let k be a field, $A = k\langle X|S\rangle = K\langle X\rangle/Id(S)$ and $<$ a monomial order on X^*, where $Id(S)$ is the ideal of $k\langle X\rangle$ generated by S. Then the following statements are equivalent:*

(i) S is a Gröbner-Shirshov basis.

(ii) $f \in Id(S) \Rightarrow \bar{f} = a\bar{s}b$ for some $s \in S$ and $a, b \in X^*$.

(iii) $Irr(S) = \{u \in X^* | u \neq a\bar{s}b, s \in S, a, b \in X^*\}$ is a basis of the algebra $A = k\langle X|S\rangle$.

The following lemma comes from [8] ([8], Lemma 4.2) which is essentially the same as Lemma 1.4.

Lemma 1.5 (Composition-Diamond Lemma) *Let k be a field, $S \subseteq K\langle X_+\rangle$ be monic, $A = k\langle X_+|S\rangle = K\langle X_+\rangle/Id(S)$ and $<$ a monomial order on X^+, where $Id(S)$ is the ideal of $k\langle X_+\rangle$ generated by S. Then the following statements are equivalent:*

(i) S is a Gröbner-Shirshov basis.

(ii) $f \in Id(S) \Rightarrow \bar{f} = a\bar{s}b$ for some $s \in S$ and $a, b \in X^*$.

(iii) $Irr(S) = \{u \in X^+ | u \neq a\bar{s}b, s \in S, a, b \in X^*\}$ is a basis of the algebra $A = k\langle X_+|S\rangle$.

Remark: Suppose that $S \subseteq K\langle X_+\rangle$ is a Gröbner-Shirshov basis. In the (iii) of Lemma 1.5, $1 \notin Irr(S)$ but not the case in the Lemma 1.4.

For convenience, we identify a relation $u = f_u$ of an algebra presented by generators and relations with the polynomial $u - f_u$ in the corresponding free algebra.

The concepts of the extension of algebras was invented by Hochschild [9] (also see [7] and [10]).

Definition 1.6 *Let k be a field, $M, B, \mathcal{R}$ k-algebras (not necessarily with 1). Then $\mathcal{R}$ is called an extension of M by B if $M^2 = 0$, M is an ideal of $\mathcal{R}$ and $\mathcal{R}/M \cong B$ as algebras.*

In [9], such an extension is called a singular extension.

2 Characterizations of extensions of algebras

Let M, B be k-algebras, $M^2 = 0$ and M a B-bimodule.

By a factor set $\{(b, b')|b, b' \in B\}$ of B in M we mean that $\{(b, b')|b, b' \in B\}$ is a subset of M such that the function (b, b') is k-bilinear.

Let $I,\ J$ be linearly ordered sets, $\{b_i|i \in I\},\ \{m_j|j \in J\}$ k-bases of B and M, respectively and $\{(b, b')|b, b' \in B\}$ a factor set of B in M. Denote

$$A = E_k(M, B, (b_p, b_q)) = k\langle(\{m_j\}, \{b_i\})_+|S\rangle$$

where $S = \{b_pb_q = [b_pb_q] + (b_p, b_q), b_pm_j = [b_pm_j], m_jb_p = [m_jb_p], m_jm_l = 0, p, q \in I, j, l \in J\}$ and for example, $[b_pb_q] = \sum_{i\in I} \alpha^i_{pq}b_i,\ \alpha^i_{pq} \in k$, is the product in B.

We order the set $(\{m_j\}_J \cup \{b_i\}_I)^+$ by the deg-lex order.

Equipping the above concepts, we have the following theorems which give characterizations of extensions of algebras.

Theorem 2.1 *Let M, B be k-algebras, $M^2 = 0$, M a B-bimodule, $\{b_i|i \in I\}$, $\{m_j|j \in J\}$ k-bases of B and M, respectively and $\{(b, b')|b, b' \in B\}$ a factor set of B in M. Let $A = E_k(M, B, (b_p, b_q)) = k\langle(\{m_j\}, \{b_i\})_+|S\rangle$, where $S = \{b_pb_q = [b_pb_q] + (b_p, b_q), b_pm_j = [b_pm_j], m_jb_p = [m_jb_p], m_jm_l = 0, p, q \in I, j, l \in J\}$. Then, S is a Gröbner-Shirshov bases for A if and only if the factor set satisfies the following condition in M: for any $p, q, r \in I$*

$$b_p(b_q, b_r) - (b_pb_q, b_r) + (b_p, b_qb_r) - (b_p, b_q)b_r = 0 \qquad (1)$$

i.e., the function (b, b') is a cocycle.

Moreover, if this is the case, $A = E_k(M, B, (b_p, b_q)) = k\langle(\{m_j\}, \{b_i\})_+|S\rangle$ is an extension of M by B in a natural way.

Proof. Suppose that (1) holds. Then the possible compositions in S are related to the following ambiguities:

$$w = b_pb_qb_r,\ m_jb_pb_q,\ b_pb_qm_j,\ m_jm_lm_n,\ m_jm_lb_p,\ b_pm_jm_l,\ b_pm_jb_q,\ m_jb_pm_l.$$

For $w = m_jm_lm_n,\ m_jm_lb_p,\ b_pm_jm_l,\ b_pm_jb_q,\ m_jb_pm_l$, by noting that $[[m_jm_l]m_n] = [m_j[m_lm_n]] = 0,\ [[m_jm_l]b_p] = [m_j[m_lb_p]] = 0,\ [[b_pm_j]m_l] = [b_p[m_jm_l]] = 0,\ [[m_jb_p]m_l] = [m_j[b_pm_l]] = 0$ and $[[b_pm_j]b_q] = [b_p[m_jb_q]]$ for any $j,l,n \in J,\ p,q \in I$, the correspondent compositions are trivial modulo S.

For $w = m_jb_pb_q$, since $[[m_jb_p]b_q] = [m_j[b_pb_q]]$ and $m_j(b_p, b_q) = 0$, we have

$$\begin{aligned} & (m_jb_p - [m_jb_p], b_pb_q - [b_pb_q] - (b_p, b_q))_w = -[m_jb_p]b_q + m_j([b_pb_q] + (b_p, b_q)) \\ \equiv\ & -[[m_jb_p]b_q] + m_j[b_pb_q] + m_j(b_p, b_q) \equiv -[[m_jb_p]b_q] + [m_j[b_pb_q]] \equiv 0 \quad mod(S, w) \end{aligned}$$

Similarly, for $w = b_pb_qm_j$, the correspondent composition is trivial modulo S.

For $w = b_pb_qb_r$, by noting that $[[b_pb_q]b_r] = [b_p[b_qb_r]]$, we have

$$\begin{aligned} & (b_pb_q - [b_pb_q] - (b_p, b_q), b_qb_r - [b_qb_r] - (b_q, b_r))_w \\ =\ & -([b_pb_q] + (b_p, b_q))b_r + b_p([b_qb_r] + (b_q, b_r)) \\ \equiv\ & -[[b_pb_q]b_r] - (b_pb_q, b_r) - (b_p, b_q)b_r + ([b_p[b_qb_r]] + (b_p, b_qb_r) + b_p(b_q, b_r)) \\ \equiv\ & 0 \quad mod(S, w) \end{aligned}$$

Thus, all compositions in S are trivial modulo S.

Conversely, if S is a Gröbner-Shirshov bases for A, then, by noting that for $w = b_pb_qb_r$, the compositions are trivial modulo S, we can easily check that the condition (1) holds in $\overline{M} = (M + Id(S))/Id(S)$. Then, by using Lemma 1.5, the algebra $\overline{M}$ is the same as M.

Moreover, assume that S is a Gröbner-Shirshov bases for A. Then, by Lemma 1.5, each element $r \in A$ can be uniquely written as $r = m + b$, where $m \in M,\ b \in B$. Hence, $A = M \oplus B$ as k-modules with the following multiplication: for any $m, m' \in M,\ b, b' \in B$,

$$(m + b) \cdot (m' + b') = mb' + bm' + (b, b') + bb'.$$

From this it follows that M is an ideal of A and $A/M \cong B$ as algebras. $\square$

Theorem 2.2 *Let $M, B, \mathcal{R}$ be k-algebras with $M^2 = 0$. If $\mathcal{R}$ is an extension of M by B and $\sigma :\ \mathcal{R}/M \to B, r_b + M \mapsto b$ an algebra isomorphism, then M is a B-bimodule in a natural way: for any $b \in B,\ m \in M$,*

$$b \cdot m = r_bm, \quad m \cdot b = mr_b \tag{2}$$

and there exists a factor set $\{(b, b')|b, b' \in B\}$ of B in M such that for any $b, b', b'' \in B$,

$$b(b', b'') - (bb', b'') + (b, b'b'') - (b, b')b'' = 0.$$

Moreover, $\mathcal{R} \cong A$ as algebras, where $A = E_k(M, B, (b_p, b_q)) = k\langle(\{m_j\}, \{b_i\})_+|S\rangle$, $\{m_j\}_J,\ \{b_i\}_I$ are any linear bases of M and B respectively and $S = \{b_pb_q = [b_pb_q] + (b_p, b_q), b_pm_j = [b_pm_j], m_jb_p = [m_jb_p], m_jm_l = 0, p, q \in I, j, l \in J\}$.

Proof. Clearly, as k-modules, $\mathcal{R}= M \oplus C$, where $C = \sum_{i\in I} kr_{b_i}$ with a basis $\{r_{b_i}|i \in I\}$. Thus, for any $b, b' \in B$, we have $r_b + r_{b'} = r_{b+b'}$ because σ is an algebra isomorphism. Since $(r_b + M)(r_{b'} + M) = r_br_{b'} + M = r_{bb'} + M$, there exists a unique $(b, b') \in M$

such that $r_b r_{b'} = r_{bb'} + (b, b')$. Then, it is easy to see that M is a B-bimodule with the module operations (2) and the function (b, b') is k-bilinear. For example, for any $b, b', b'' \in B, m \in M$, $(bb')\cdot m = r_{bb'}m = (r_b r_{b'} - (b, b'))m = (r_b r_{b'})m = r_b(r_{b'}m) = b\cdot(b'\cdot m)$. Also, since $r_b(r_{b'} + r_{b''}) = r_b r_{b'} + r_b r_{b''} = r_b r_{b'+b''}$, we have $(b, b' + b'') = (b, b') + (b, b'')$.

Moreover, by noting that $(r_b r_{b'})r_{b''} = r_b(r_{b'}r_{b''})$, we know that the factor set $\{(b, b')|b, b' \in B\}$ satisfies

$$b(b', b'') - (bb', b'') + (b, b'b'') - (b, b')b'' = 0.$$

Now, we prove that $\mathcal{R} \cong A$ as algebras. For any $m, m' \in M,\ c, c' \in C$,

$$(m + c) \cdot (m' + c') = mc' + cm' + (c, c') + cc'$$

where $(c, c') = \sum_{p,q} \alpha_p \alpha'_q (b_p, b_q)$ if $c = \sum_p \alpha_p r_{b_p}$, $c' = \sum_q \alpha'_q r_{b_q}$. Then, by the proof of Theorem 2.1, it is easy to see that $\tau :\ \mathcal{R} \to A$ by $\tau(m_j) = m_j$, $\tau(r_{b_i}) = b_i$ is an algebra isomorphism. □

By Theorem 2.1 and Theorem 2.2, we have the following theorem.

Theorem 2.3 *Let $M, B, \mathcal{R}$ be k-algebras with $M^2 = 0$. Then $\mathcal{R}$ is an extension of M by B if and only if M is a B-bimodule and there exists a factor set $\{(b, b')|b, b' \in B\}$ of B in M such that for any $b, b', b'' \in B$,*

$$b(b', b'') - (bb', b'') + (b, b'b'') - (b, b')b'' = 0$$

and $\mathcal{R} \cong A = E_k(M, B, (b_p, b_q)) = k\langle(\{m_j\}, \{b_i\})_+|S\rangle$, where $\{b_i\}_I$ and $\{m_j\}_J$ are any linear bases of B and M respectively, $S = \{b_p b_q = [b_p b_q] + (b_p, b_q), b_p m_j = [b_p m_j], m_j b_p = [m_j b_p], m_j m_l = 0, p, q \in I, j, l \in J\}$.

Now, we consider the general case when the algebra B is presented by generators and relations.

Let M, B be k-algebras, $M^2 = 0$, M a B-bimodule, $B = k\langle X_+|R\rangle$, where R is a Gröbner-Shirshov bases for B with the deg-lex order $<_B$ on X^+.

For convenience, we can assume that R is a minimal Gröbner-Shirshov bases in a sense that the leading monomials are not contained each other as subwords, in particular, they are pairwise different. Let $R = \{u - f_u|u \in \Lambda\}$, where u is the leading term of the polynomial $h_u = u - f_u$ in $k\langle X_+\rangle$. Let

$$A = E_k(M, X, (u)) = k\langle(\{m_j\} \cup X)_+|S_1\rangle$$

where $\{m_j\}_J$ is a basis of M, $S_1 = \{u = f_u + (u), u \in \Lambda,\ xm_j = [xm_j], m_j x = [m_j x],\ m_j m_l = 0, x \in X, j, l \in J\}$ and $\{(u)|u \in \Lambda\} \subseteq M$. For convenience, we also call $\{(u)|u \in \Lambda\}$ a factor set of B in M.

We order the set $(\{m_j\}_J \cup X)^+$ also by the deg-lex order which extends $<_B$ and satisfies $x > m_j$ for any $x \in X$ and $j \in J$.

In S_1, the possible compositions are related to the following ambiguities:

$$w_1 = w,\ m_j u,\ u m_j,\ m_j m_l m_n,\ m_j m_l x,\ x m_j m_l,\ x m_j x',\ m_j x m_l$$

where w is an ambiguity appeared in R.

For $w_1 = m_jm_lm_n,\ m_jm_lx,\ xm_jm_l,\ xm_jx',\ m_jxm_l$, by noting that $[[m_jm_l]m_n] = [m_j[m_lm_n]] = 0,\ [[m_jm_l]x] = [m_j[m_lx]] = 0,\ [[xm_j]m_l] = [x[m_jm_l]] = 0,\ [[m_jx]m_l] = [m_j[xm_l]] = 0$ and $[[xm_j]x'] = [x[m_jx']]$ for any $j, l, n \in J,\ x, x' \in X$, the corresponding compositions are trivial modulo S_1.

For $w_1 = m_ju,\ u = x_1 \cdots x_t$, we have

$$\begin{aligned} & (m_jx_1 - [m_jx_1], u - (f_u + (u)))_{w_1} = -[m_jx_1]x_2 \cdots x_t + m_j(f_u + (u)) \\ \equiv\ & -[\cdots[[m_jx_1]x_2] \cdots x_t] + m_jf_u + m_j(u) \equiv -[m_ju] + [m_jf_u] \equiv 0 \ \ mod(S_1, w_1) \end{aligned}$$

Similarly, for $w_1 = um_j$, we have $(u - (f_u + (u)), xm_j - [xm_j])_{w_1} \equiv 0 \ \ mod(S_1, w_1)$.

Since R is a minimal Gröbner-Shirshov bases, all compositions in R are only intersection ones.

Now, for $w_1 = w = u_1c = du_2,\ u_1, u_2 \in \Lambda,\ c, d \in X^+$, we have

$$\begin{aligned} & (u_1 - (f_{u_1} + (u_1)), u_2 - (f_{u_2} + (u_2)))_{w_1} = -(f_{u_1} + (u_1))c + d(f_{u_2} + (u_2)) \\ =\ & (df_{u_2} - f_{u_1}c) + (d(u_2) - (u_1)c) \end{aligned}$$

Since the composition $(h_{u_1}, h_{u_2})_{w_1}$ in R is trivial modulo R, it follows that in $k\langle X_+\rangle$,

$$(h_{u_1}, h_{u_2})_{w_1} = df_{u_2} - f_{u_1}c = \sum \alpha_i a_i(u'_i - f_{u'_i})b_i \tag{3}$$

where each $\alpha_i \in k,\ a_i, b_i \in X^*,\ u'_i \in \Lambda$ and $\overline{a_i(u'_i - f_{u'_i})b_i} = \overline{a_iu'_ib_i} < w_1$. Therefore, in $k\langle(\{m_j\} \cup X)_+\rangle$, we have

$$\begin{aligned} & (u_1 - (f_{u_1} + (u_1)), u_2 - (f_{u_2} + (u_2)))_{w_1} = (df_{u_2} - f_{u_1}c) + (d(u_2) - (u_1)c) \\ =\ & \sum \alpha_i a_i(u'_i - f_{u'_i} - (u'_i))b_i + \sum \alpha_i a_i(u'_i)b_i - ((u_1)c - d(u_2)) \end{aligned}$$

where $\overline{a_i(u'_i - f_{u'_i} - (u'_i))b_i} = \overline{a_iu'_ib_i} < w_1$. From this it follows that

$$(u_1 - (f_{u_1} + (u_1)), u_2 - (f_{u_2} + (u_2)))_{w_1} \equiv g_{(u_1,u_2)_w}(u) \ \ mod(S_1, w_1) \tag{4}$$

where $g_{(u_1,u_2)_w}(u) = \sum \alpha_i a_i(u'_i)b_i - ((u_1)c - d(u_2)) \in M$ is a function of $\{(u) | u \in \Lambda\}$.

In fact, by (3), we have an algorithm to find the function $g_{(u_1,u_2)_w}$.

Therefore, we have the following theorem.

Theorem 2.4 *Let $B = k\langle X_+ | R\rangle$, where $R = \{u - f_u | u \in \Lambda\}$ is a minimal Gröbner-Shirshov bases for B and u the leading term of the polynomial $h_u = u - f_u$ in $k\langle X_+\rangle$. Let M be a k-module, $M^2 = 0$, M a B-bimodule, $\{m_j\}_J$ a k-basis of M and $\{(u) | u \in \Lambda\}$ a factor set of B in M. Let $A = E_k(M, X, (u)) = k\langle(\{m_j\}_J \cup X)_+ | S_1\rangle$ where $S_1 = \{u = f_u + (u), xm_j = [xm_j], m_jx = [m_jx], m_jm_l = 0, u \in \Lambda, x \in X, j, l \in J\}$. Then, S_1 is a Gröbner-Shirshov bases for A if and only if*

$$\{g_{(u_1,u_2)_w}(u) | (h_{u_1}, h_{u_2})_w \ \textit{is a composition in}\ R\} = 0 \tag{5}$$

where $g_{(u_1,u_2)_w}(u)$ is defined by (4). Moreover, if this is the case, A is an extension of M by B in a natural way.

Proof. Assume that S_1 is a Gröbner-Shirshov bases for A. Then, by the previous statements, for any composition $(U_1, U_2)_w$ in R, $g_{(u_1,u_2)_w} = 0$ in $\overline{M} = (M + Id(S))/Id(S)$. Then, by using Lemma 1.5, the algebra $\overline{M}$ is the same as M. So, the (5) holds. Conversely, if (5) holds, then it is clear that S_1 is a Gröbner-Shirshov bases for A.

We need only to prove that A is an extension of M by B in a natural way if (5) holds. By Lemma 1.5, each element $r \in A$ can be uniquely written as $r = m + b$, where $m \in M,\ b \in B$ is R-irreducible. Hence, $A = M \oplus B$ as k-modules with the following multiplication: for any $m, m' \in M,\ b, b' \in B$,

$$(m + b) \cdot (m' + b') = mb' + bm' + (u_{b,b'}) + bb'$$

where $bb' \equiv [bb']\ mod(R),\ [bb']$ is R-irreducible and $(u_{b,b'}) \in M$ is a function of $\{(u)|u \in \Lambda\}$. From this it follows that M is an ideal of A and $A/M \cong B$ as algebras. □

Theorem 2.5 *Let $B = k\langle X_+|R\rangle$, where $R = \{u - f_u|u \in \Lambda\}$ is a minimal Gröbner-Shirshov bases for B and u the leading term of the polynomial $h_u = u - f_u$ in $k\langle X_+\rangle$. Let $M, \mathcal{R}$ be k-algebras with $M^2 = 0$. If $\mathcal{R}$ is an extension of M by B and $\sigma : \mathcal{R}/M \to B,\ r_x + M \mapsto x$ an algebra isomorphism, then M is a B-bimodule with a natural way: for any $x \in X,\ m \in M$,*

$$x \cdot m = r_x m, \quad m \cdot x = m r_x$$

and there exists a factor set $\{(u)|u \in \Lambda\}$ of B in M such that

$$\{g_{(u_1,u_2)_w}(u)|(h_{u_1}, h_{u_2})_w \text{ is a composition in } R\} = 0$$

where $g_{(u_1,u_2)_w}(u)$ is defined by (4). Moreover, $\mathcal{R} \cong A$ as algebras, where $A = E_k(M, X, (u)) = k\langle(\{m_j\}_J \cup \{X\})_+|S_1\rangle$, $\{m_j\}_J$ a k-basis of M and $S_1 = \{u = f_u + (u), xm_j = [xm_j], m_j x = [m_j x], m_j m_l = 0, u \in \Lambda, x \in X, j, l \in J\}$.

Proof. It is clear that M is a B-bimodule under the given operations.

For any $u \in \Lambda$, since $r_u + M = r_{f_u} + M$, there exists a unique $(u) \in M$ such that $r_u = r_{f_u} + (u)$. By noting that each composition $(h_{u_1}, h_{u_2})_w$ in R, $(h_{u_1}, h_{u_2})_w \equiv 0\ mod(R, w)$, we have that $g_{(u_1,u_2)_w}(u) = 0$ in M by (3).

Now, by using the proof of Theorem 2.4, S_1 is a Gröbner-Shirshov bases for the algebra A and then the result follows. □

The following theorem, which gives a complete characterization of an algebra to be an extension of M by B, follows from Theorem 2.4 and Theorem 2.5.

Theorem 2.6 *Let $M, B, \mathcal{R}$ be k-algebras with $M^2 = 0,\ B = k\langle X_+|R\rangle$, where $R = \{u - f_u|u \in \Lambda\}$ is a minimal Gröbner-Shirshov bases for B and u the leading term of the polynomial $h_u = u - f_u$ in $k\langle X_+\rangle$. Then $\mathcal{R}$ is an extension of M by B if and only if M is a B-bimodule and there exists a factor set $\{(u)|u \in \Lambda\}$ of B in M such that*

$$\{g_{(u_1,u_2)_w}(u)|(h_{u_1}, h_{u_2})_w \text{ is a composition in } R\} = 0$$

and $\mathcal{R} \cong E_k(M, X, (u)) = k\langle(\{m_j\}_J \cup \{X\})_+|S_1\rangle$, where $g_{(u_1,u_2)_w}(u)$ is defined by (4), $\{m_j\}_J$ a k-basis of M and $S_1 = \{u = f_u + (u), xm_j = [xm_j], m_j x = [m_j x], m_j m_l = 0, u \in \Lambda, x \in X, j, l \in J\}$.

3 Applications

Let $M, B, \mathcal{R}$ be k-algebras with $M^2 = 0$. The previous theorems give an answer to how to find the conditions which makes $\mathcal{R}$ to be an extension of M by B. We call the condition (4) the extension condition of M by B. In fact, for the extension condition, it is essential to find the function $g_{(u_1,u_2)_w}(u)$, which can be obtained by algorithm (3).

As results, by using the extension conditions, let us give some examples. We give the characterization of the extension of M by B when the B is cyclic algebra, free commutative algebra, universal envelope of the free Lie algebra and Grassman algebra respectively.

Theorem 3.1 *Let $M, B, \mathcal{R}$ be k-algebras with $M^2 = 0$ and $B = k\langle\{x\}_+|x^n = f(x)\rangle$ a cyclic algebra, where n is a natural number and $f(x)$ is a polynomial of degree less than n such that $f(0) = 0$. Then, $\mathcal{R}$ is isomorphic to an extension of M by B if and only if M is a B-bimodule, there exists an $m \in M$ such that $mx = xm$ and $\mathcal{R} \cong E_k(M, x, m) = k\langle(\{m_j\}_J \cup \{x\})_+|S\rangle$, where $\{m_j\}_J$ is any k-basis of M and $S = \{x^n = f(x) + m,\ xm_j = [xm_j], m_jx = [m_jx], m_jm_l = 0, j, l \in J\}$.*

Proof. Clearly, $R = \{x^n = f(x)\}$ is a Gröbner-Shirshov bases for B. We need only to consider the composition in S: $(x^n - f(x) - m, x^n - f(x) - m)_w,\ w = x^{n+1}$. Thus, we obtain the extension condition: $mx = xm$. Now, by Theorem 2.6, the result follows. $\square$

Theorem 3.2 *Let $M, B, \mathcal{R}$ be k-algebras with $M^2 = 0$. Let $X = \{x_i|i \in I\}$, I a well ordered set and $B = k\langle X_+|x_px_q = x_qx_p,\ p > q,\ p, q \in I\rangle$ the free commutative algebra generated by X. Then, $\mathcal{R}$ is isomorphic to an extension of M by B if and only if M is a B-bimodule, there exists a factor set $\{(x_p, x_q)|p > q,\ p, q \in I\}$ of B in M such that for any $p, q, r \in I,\ p > q > r$,*

$$(x_q, x_r)x_p - x_p(x_q, x_r) + x_q(x_p, x_r) - (x_p, x_r)x_q + (x_p, x_q)x_r - x_r(x_p, x_q) = 0 \qquad (6)$$

and $\mathcal{R} \cong E_k(M, X, (x_p, x_q)) = k\langle(\{m_j\}_J \cup X)_+|S\rangle$, where $\{m_j\}_J$ is any k-basis of M and $S = \{x_px_q = x_qx_p + (x_p, x_q),\ x_pm_j = [x_pm_j], m_jx_p = [m_jx_p], m_jm_l = 0, p > q,\ p, q \in I, j, l \in J\}$.

Proof. Let $R = \{x_px_q = x_qx_p|p > q,\ p, q \in I\}$. Then, for the deg-lex order on X^+, R is clearly a Gröbner-Shirshov bases for B and only one kind of compositions are in R, i.e., $(x_px_q - x_qx_p, x_qx_r - x_rx_q)_w,\ w = x_px_qx_r,\ p, q, r \in I,\ p > q > r$. Then, in S, by calculating the composition $(x_px_q - x_qx_p - (x_p, x_q), x_qx_r - x_rx_q - (x_q, x_r))_w,\ w = x_px_qx_r,\ p, q, r \in I,\ p > q > r$, we obtain the extension condition (6). Now, by Theorem 2.6, the result follows. $\square$

The following theorem is a generation of Theorem 3.2.

Theorem 3.3 *Let $M, B, \mathcal{R}$ be k-algebras with $M^2 = 0$. Let $X = \{x_i|i \in I\}$, I be a well ordered set, $L = Lie_k(X|[x_px_q] = \sum_{i\in I}\alpha^i_{pq}x_i,\ p, q \in I)$ a Lie algebra and $B = U(L) = k\langle X_+|R\rangle$ the universal envelope of the Lie algebra L, where each $\alpha^i_{pq} \in k$ and $R = \{x_px_q = x_qx_p + [x_px_q]|p > q,\ p, q \in I\}$. Then, $\mathcal{R}$ is isomorphic to an extension of M*

by B if and only if M is a B-bimodule, there exists a factor set $\{(x_p,x_q)|p,q\in I\}$ of B in M such that for any $p,q,r\in I,\ p>q>r$,

$$\begin{aligned}&(x_q,x_r)x_p+x_q(x_p,x_r)+(x_p,x_q)x_r+([x_qx_r],x_p)+(x_q,[x_px_r])+([x_px_q],x_r)\\&-(x_p,x_r)x_q-x_p(x_q,x_r)-x_r(x_p,x_q)=0\end{aligned}$$

and $\mathcal{R}\cong E_k(M,X,(x_p,x_q))=k\langle(\{m_j\}_J\cup X)_+|S\rangle$, where $(x_p,x_q)=-(x_q,x_p),\ (x_p,x_p)=0$ for any $p,q\in I$, $\{m_j\}_J$ is any k-basis of M and $S=\{x_px_q=x_qx_p+[x_px_q]+(x_p,x_q),\ x_pm_j=[x_pm_j],m_jx_p=[m_jx_p],m_jm_l=0,p>q,\ p,q\in I,j,l\in J\}$.

Proof. By [4], for the deg-lex order on X^+, R is a Gröbner-Shirshov bases for B and only one kind of compositions are in R, i.e., $(x_px_q-x_qx_p-[x_px_q],x_qx_r-x_rx_q-[x_qx_r])_w,\ w=x_px_qx_r,\ p,q,r\in I,\ p>q>r$. Since, in S,

$$\begin{aligned}&(x_px_q-x_qx_p-[x_px_q]-(x_p,x_q),x_qx_r-x_rx_q-[x_qx_r]-(x_q,x_r))_w\\=\ &-(x_qx_px_r+[x_px_q]x_r+(x_p,x_q)x_r)+(x_px_rx_q+x_p[x_qx_r]+x_p(x_q,x_r))\\\equiv\ &-(x_q(x_rx_p+[x_px_r]+(x_p,x_r))+[x_px_q]x_r+(x_p,x_q)x_r)\\&+((x_rx_p+[x_px_r]+(x_p,x_r))x_q+x_p[x_qx_r]+x_p(x_q,x_r))\\\equiv\ &-x_qx_rx_p-(x_q[x_px_r]+[x_px_q]x_r)-(x_q(x_p,x_r)+(x_p,x_q)x_r)\\&+x_rx_px_q+[x_px_r]x_q+x_p[x_qx_r]+(x_p,x_r)x_q+x_p(x_q,x_r)\\\equiv\ &-(x_rx_q+[x_qx_r]+(x_q,x_r))x_p+x_r(x_px_q+[x_px_q]+(x_p,x_q))\\&-(x_q[x_px_r]+[x_px_q]x_r)-(x_q(x_p,x_r)+(x_p,x_q)x_r)\\&+[x_px_r]x_q+x_p[x_qx_r]+(x_p,x_r)x_q+x_p(x_q,x_r)\\\equiv\ &-(x_q,x_r)x_p-x_q(x_p,x_r)-(x_p,x_q)x_r+(x_p,x_r)x_q+x_p(x_q,x_r)+x_r(x_p,x_q)\\&-[x_qx_r]x_p-x_q[x_px_r]-[x_px_q]x_r+[x_px_r]x_q+x_p[x_qx_r]+x_r[x_px_q]\end{aligned}$$

and by using Jacobi identity $[[x_px_r]x_q]+[x_p[x_qx_r]]+[x_r[x_px_q]]=0$,

$$\begin{aligned}&[x_px_r]x_q+x_p[x_qx_r]+x_r[x_px_q]\\\equiv\ &(x_q[x_px_r]+[[x_px_r]x_q]+([x_px_r],x_q))+([x_qx_r]x_p+[x_p[x_qx_r]]+(x_p,[x_qx_r]))\\&+([x_px_q]x_r+[x_r[x_px_q]]+(x_r,[x_px_q]))\\\equiv\ &[[x_px_r]x_q]+[x_p[x_qx_r]]+[x_r[x_px_q]]+[x_qx_r]x_p+(x_q[x_px_r]+[x_px_q]x_r\\&+(x_p,[x_qx_r])+(x_r,[x_px_q])+([x_px_r],x_q))\\\equiv\ &[x_qx_r]x_p+(x_q[x_px_r]+[x_px_q]x_r+(x_p,[x_qx_r])+(x_r,[x_px_q])+([x_px_r],x_q))\end{aligned}$$

we have the extension condition mentioned in the theorem. Now, by Theorem 2.6, the result follows. □

Theorem 3.4 *Let $M,B,\mathcal{R}$ be k-algebras with $M^2=0$. Let $X=\{x_i|i\in I\},\ I$ be a well ordered set, $B=k\langle X_+|R\rangle$ the Grassman algebra, where $R=\{x_q^2=0,\ x_px_q=-x_qx_p|p>q,\ p,q\in I\}$. Then, $\mathcal{R}$ is isomorphic to an extension of M by B if and only if M is a B-bimodule, there exists a factor set $\{(x_p,x_q)|p\geq q,\ p,q\in I\}$ of B in M such that for any $p,q,r\in I,\ p>q>r$,*

$$\begin{aligned}&(x_q,x_q)x_r-x_r(x_q,x_q)+(x_q,x_r)x_q-x_q(x_q,x_r)=0\\&(x_r,x_r)x_q-x_q(x_r,x_r)+(x_q,x_r)x_r-x_r(x_q,x_r)=0\\&(x_q,x_r)x_p+(x_p,x_r)x_q+(x_p,x_q)x_r-x_q(x_p,x_r)-x_p(x_q,x_r)-x_r(x_p,x_q)=0\end{aligned}$$

and $\mathcal{R} \cong E_k(M, X, (x_p, x_q)) = k\langle(\{m_j\}_J \cup X)_+|S\rangle$*, where* $\{m_j\}_J$ *is any* k*-basis of* M *and* $S = \{x_q^2 = (x_q, x_q),\ x_p x_q = -x_q x_p + (x_p, x_q),\ x_q m_j = [x_q m_j], m_j x_q = [m_j x_q], m_j m_l = 0, p > q,\ p, q \in I, j, l \in J\}$.

Proof. By [4], for the deg-lex order on X^+, R is a Gröbner-Shirshov bases for B and the possible compositions in R are related to the following ambiguities: $w_1 = x_q^2 x_r,\ w_2 = x_q x_r^2,\ w_3 = x_p x_q x_r,\ p, q, r \in I,\ p > q > r$. Then, in S, by calculating the corresponding compositions: $(x_q^2 - (x_q, x_q), x_q x_r + x_r x_q - (x_q, x_r))_{w_1},\ (x_q x_r + x_r x_q - (x_q, x_r), x_r^2 - (x_r, x_r))_{w_2},\ ((x_p x_q + x_q x_p - (x_p, x_q), x_q x_r + x_r x_q - (x_q, x_r))_{w_3}$, respectively, we obtain the extension conditions mentioned in the theorem. Now, by Theorem 2.6, the result follows. □

Acknowledgement: The author would like to express his deepest gratitude to Professor L. A. Bokut for his kind guidance, useful discussions and enthusiastic encouragement when the author was visiting Sobolev Institute of Mathematics.

References

[1] G. M. Bergman, The diamond lemma for ring theory, *Adv. in Math.*, 29, 178-218(1978).

[2] L.A. Bokut, Unsolvability of the word problem, and subalgebras of finitely presented Lie algebras, *Izv. Akad. Nauk. SSSR Ser. Mat.*, 36, 1173-1219(1972).

[3] L.A. Bokut, Imbeddings into simple associative algebras, *Algebra i Logika*, 15, 117-142(1976).

[4] L.A.Bokut, Y.Fong and W.F.Ke, Gröbner Shirshov bases and composition lemma for associative conformal algebras: an example, *Contemp. Math.* 264 (2000), 63-90.

[5] B. Buchberger, An algorithm for finding a basis for the residue class ring of a zero-dimensional polynomial ideal [in German], Ph.D. thesis, University of Innsbruck, Austria, (1965).

[6] B. Buchberger, An algorithmical criteria for the solvability of algebraic systems of equations[in German], *Aequationes Math.*, 4, 374-383(1970).

[7] Henri Cartan and Samuel Eilenberg, Homological Algebra, Princeton University Press, 1956.

[8] Yuqun Chen and Jianjun Qiu, Grobner-Shirshov basis for free product of algebras and beyond, *SEA. Bull. Math.*, 30, 811-826(2006).

[9] G. Hochschild, On the cohomology groups of an associative algebra, Ann. Math., 46(1945), 58-67.

[10] Saunders Maclane, Homology, Springer-Verlag, 1963.

[11] A.I. Shirshov, Some algorithmic problem for Lie algebras, *Sibirsk. Mat. Z.*, 3(1962), 292-296(in Russian); English translation in SIGSAM Bull., 33(2), 3-6(1999).